

Bordism of Unoriented Surfaces in 4-Space

J. Scott Carter
University of South Alabama
Mobile, AL 36688
carter@mathstat.usouthal.edu

Seiichi Kamada
Osaka City University
Osaka 558-8585, JAPAN
kamada@sci.osaka-cu.ac.jp

Masahico Saito
University of South Florida
Tampa, FL 33620
saito@math.usf.edu

Shin Satoh
Kyoto University
Kyoto 606-8502, JAPAN
satoh@kurims.kyoto-u.ac.jp

February 1, 2008

Abstract

The group of bordism classes of unoriented surfaces in 4-space is determined. The bordism classes are characterized by normal Euler numbers, double linking numbers, and triple linking numbers.

1 Introduction

B. J. Sanderson [9, 10] studied the group $L_{m,n}$ of bordism classes of ‘oriented’ closed $(m-2)$ -manifolds of n components in $\mathbf{R}^m$. He showed that $L_{m,n}$ is isomorphic to the homotopy group $\pi_m(\bigvee_{i=1}^{n-1} S^2)$; in particular, the bordism group $L_{m,n}$ for $m = 4$ is given as follows.

Theorem 1.1 (Sanderson)

$$L_{4,n} \cong \underbrace{(\mathbf{Z}_2 \oplus \dots \oplus \mathbf{Z}_2)}_{\frac{n(n-1)}{2}} \oplus \underbrace{(\mathbf{Z} \oplus \dots \oplus \mathbf{Z})}_{\frac{n(n-1)(n-2)}{3}}.$$

In particular, we have $L_{4,1} \cong \{0\}$, $L_{4,2} \cong \mathbf{Z}_2$, and $L_{4,3} \cong \mathbf{Z}_2^3 \oplus \mathbf{Z}^2$.

Similarly, there is a group of bordism classes of ‘unoriented’ closed $(m-2)$ -manifolds of n components in $\mathbf{R}^m$. We denote the group by $UL_{m,n}$. The aim of this paper is to determine the bordism group $UL_{m,n}$ for $m = 4$ via purely geometric techniques.

An n -component surface-link $F = F_1 \cup \dots \cup F_n$ is a disjoint union of closed surfaces embedded in $\mathbf{R}^4$ (smoothly, or PL and locally flatly) such that each component F_i , $i = 1, \dots, n$, is an orientable or non-orientable embedded surface, which may or may not be connected. Two n -component surface-links F and F' are *unorientedly bordant* if there is a compact 3-manifold $W = \bigcup_{i=1}^n W_i$ properly embedded in $\mathbf{R}^4 \times [0, 1]$ such that $\partial W_i = F_i \times \{0\} \cup F'_i \times \{1\}$ for $i = 1, \dots, n$. In this paper, $F \simeq_B F'$ means that F and F' are

unorientedly bordant, and $F \cong_A F'$ means that they are ambient isotopic in $\mathbf{R}^4$. The unoriented bordism classes of n -component surface-links form an abelian group $UL_{4,n}$ such that the sum $[F] + [F']$ is defined to be the class $[F \amalg F']$ of the split union $F \amalg F'$. The identity is represented by the empty $F = \emptyset$ and the inverse $-[F]$ is represented by the mirror image of F . The following is our main theorem.

Theorem 1.2

$$UL_{4,n} \cong \underbrace{(\mathbf{Z} \oplus \dots \oplus \mathbf{Z})}_n \oplus \underbrace{(\mathbf{Z}_4 \oplus \dots \oplus \mathbf{Z}_4)}_{\frac{n(n-1)}{2}} \oplus \underbrace{(\mathbf{Z}_2 \oplus \dots \oplus \mathbf{Z}_2)}_{\frac{n(n-1)(n-2)}{3}}.$$

In particular, we have $UL_{4,1} \cong \mathbf{Z}$, $UL_{4,2} \cong \mathbf{Z}^2 \oplus \mathbf{Z}_4$, and $UL_{4,3} \cong \mathbf{Z}^3 \oplus \mathbf{Z}_4^3 \oplus \mathbf{Z}_2^2$.

This paper is organized as follows. In §2, we give definitions of three kinds of unoriented bordism invariants — normal Euler numbers, double linkings, and triple linkings — by the projection method. In §3, we study 1-component surface-links. In §4, we introduce a family of surface-links which are called *necklaces*. §5 is devoted to study crossing changes that produce necklaces. In §6, we prove Theorem 1.2.

2 Unoriented Bordism Invariants

All of our bordism invariants will be defined using the diagram of a knotted or linked surface. We begin by recalling this notion. Consider a surface-link F . We may assume that the restriction $\pi|_F : F \rightarrow \mathbf{R}^3$ of a projection $\pi : \mathbf{R}^4 \rightarrow \mathbf{R}^3$ is a generic map, that is, the singularity set of the image $\pi(F) \subset \mathbf{R}^3$ consists of double points and isolated branch/triple points. See Figure 1. The closure of the self-intersection set on $\pi(F)$ is regarded as a union of immersed arcs and loops, which we call double curves. Branch points (or Whitney umbrella points) occur at the end of the double curves, and triple points occur when double curves intersect.




Figure 1: Generic intersections of surfaces in 3-space

A *surface diagram* of F is the image $\pi(F)$ equipped with over/under-information along each double curve with respect to the projection direction. To indicate such over/under-information, we remove a neighborhood of a double curve on the sheet (the *lower sheet*) which lies lower than the other sheet (the *upper sheet*). See Figure 2. Notice that the removal of this neighborhood is merely a convention in depicting illustrations. In particular, we still

speak of “double curves” and triple points, and we locally parametrize the surfaces using immersions.

There are seven kinds of local moves on surface diagrams, called *Roseman moves*, analogues of Reidemeister moves for classical knots, which are sufficient to relate diagrams of ambient isotopic surface-links (cf. [3, 4, 8]). Specifically, two diagrams represent ambiently isotopic surface-links if and only if one can be obtained from the other by means a finite sequence of moves taken from the list of Roseman moves. We remark that the Roseman moves are used here only in relation their affect on the cobordism invariants.




Figure 2: The broken surface diagrams at intersection points

For a surface-knot K (that is a connected closed surface embedded in $\mathbf{R}^4$), H. Whitney defined the *normal Euler number* $e(K)$ of K to be the Euler number of a tubular neighborhood of K in $\mathbf{R}^4$ considered as a 2-plane bundle (cf. [7, 12]). It is known that

- (i) $e(K) = 0$ if K is orientable,
- (ii) (Whitney’s congruence) $e(K) \equiv 2\chi(K) \pmod{4}$, and
- (iii) (Whitney-Massey theorem) $|e(K)| \leq 4 - 2\chi(K)$,

where $\chi(K)$ denotes the Euler characteristic of K (cf. [6, 7]). For a 1-component surface-link F , we define the normal Euler number $e(F)$ of F to be the sum of $e(K)$ for the connected components K of F . The normal Euler number $e(F)$ can be calculated by use of a projection of F in $\mathbf{R}^3$; it is equal to the number of positive type branch points (see Figure 2) minus that of negative type ones, [2].

Let $F = F_1 \cup F_2$ be a 2-component surface-link and D a surface diagram of F . A double curve of D is said to be of *type* (i, j) if the upper sheet belongs to F_i and the lower belongs to F_j , where $i, j \in \{1, 2\}$. If a double curve is an immersed arc, then its endpoints are branch points, and hence, the type is $(1, 1)$ or $(2, 2)$. Let $C = c_1 \cup \dots \cup c_m$ be the set of the double curves of type $(1, 2)$ on the surface diagram D . Each double curve c_i is an immersed loop in $\mathbf{R}^3$.

We take a 2-disk B^2 and a union of intervals X in $\mathbf{R}^2$ as follows:

- (i) $B^2 = \{(x, y) | x^2 + y^2 \leq 1\}$, and
- (ii) $X = \{(x, y) | -1 \leq x \leq 1, y = 0\} \cup \{(x, y) | x = 0, -1 \leq y \leq 1\}$.

For a regular neighborhood $N(c_i)$ of c_i in $\mathbf{R}^3$, the pair $(N(c_i), D \cap N(c_i))$ is regarded as the image of an immersion, say φ , of one of the following manifold pairs:

- (i) $(B^2, X) \times [0, 1] / (x, 0) \sim (x, 1)$ for $x \in B^2$, and

(ii) $(B^2, X) \times [0, 1]/(x, 0) \sim (-x, 1)$ for $x \in B^2$.

Let c'_i be a loop or a pair of loops immersed in $N(c_i)$ such that $c'_i = \varphi(\{z, -z\} \times [0, 1]/\sim)$ for some $z \in B^2 \setminus X$. We give orientations to c_i and c'_i such that $[c'_i] = 2[c_i] \in H_1(N(c_i); \mathbf{Z})$. See Figure 3. We put $C' = c'_1 \cup \dots \cup c'_m$. Since C and C' are mutually disjoint 1-cycles in $\mathbf{R}^3$, the linking number $\text{Lk}(C, C')$ between C and C' is defined; let d be a 2-cycle in $\mathbf{R}^3$ with $\partial d = C$ such that d and C' intersect transversely. Then $\text{Lk}(C, C')$ is the algebraic intersection number of C' with d . This number is well-defined modulo 4; it does not depend on a choice of $z \in B^2 \setminus X$ and an orientation of c_i for each i . Furthermore, its congruence class modulo 4 remains unchanged under the Roseman moves. Hence, the mod 4 reduction of $\text{Lk}(C, C')$ is an ambient isotopy invariant of $F = F_1 \cup F_2$.




Figure 3: The double curve push-offs

Definition 2.1 The *double linking number* between F_1 and F_2 , denoted by $d(F_1, F_2)$, is a value in $\mathbf{Z}_4 = \mathbf{Z}/4\mathbf{Z} = \{0, 1, 2, 3\}$ that is the linking number $\text{Lk}(C, C')$ modulo 4.

It is proved later that the double linking number is asymmetric; $d(F_1, F_2) = -d(F_2, F_1)$.

Let $F = F_1 \cup F_2 \cup F_3$ be a 3-component surface-link. At a triple point on a surface diagram of F , there are three sheets called *top*, *middle*, and *bottom* with respect to the projection direction. A triple point is of *type* (i, j, k) if the top sheet comes from F_i , the middle comes from F_j , and the bottom comes from F_k , where $i, j, k \in \{1, 2, 3\}$. Let $N(i, j, k)$ denote the number of the triple points of type (i, j, k) . The mod 2 reduction of $N(i, j, k)$ is preserved under Roseman moves and hence an ambient isotopy invariant of F provided $i \neq j$ and $j \neq k$ (possibly $i = k$), [11].

Definition 2.2 The *triple linking number* among F_i, F_j , and F_k , denoted by $t(F_i, F_j, F_k)$, is a value in $\mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z} = \{0, 1\}$ that is the number $N(i, j, k)$ modulo 2 provided $i \neq j$ and $j \neq k$.

Lemma 2.3 The ambient isotopy invariants e, d and t are unoriented bordism invariants.

Proof. We take surface diagrams D and D' of surface-links F and F' respectively. If F and F' are unorientably bordant, then D' is obtained from D by a finite sequence of moves from the following list:

- an ambient isotopy of $\mathbf{R}^3$,
- a Roseman move,
- adding or deleting embedded 2-spheres in $\mathbf{R}^3$ which are disjoint from D ,
- a 1-handle surgery on D in $\mathbf{R}^3$ whose core is a simple arc γ with $\gamma \cap D = \partial\gamma$ (see Figure 4), and
- a 2-handle surgery on D in $\mathbf{R}^3$ whose core is a simple 2-disk δ with $\delta \cap D = \partial\delta$.

Recall that Roseman moves do not change the invariants e, d and t . The other deformations listed above do not change the singularity set of the diagram. Hence e, d and t are unoriented bordism invariants. ■




Figure 4: Attaching 1 and 2-handles

3 1-Component Surface-Links

A projective plane embedded in $\mathbf{R}^4$ is *standard* if it has a surface diagram as shown in Figure 5. A non-orientable surface-knot is said to be *trivial* if it is a connected sum of some standard projective planes in $\mathbf{R}^4$. Two trivial non-orientable surface-knots F and F' are ambient isotopic if and only if $e(F) = e(F')$ and $\chi(F) = \chi(F')$. The following lemma is folklore.




Figure 5: The positive and negative projective planes

Lemma 3.1 *Two 1-component surface-links F and F' are unorientedly bordant if and only if $e(F) = e(F')$.*

Proof. The only if part is obvious. We prove the if part. It is known that any non-orientable surface-link is transformed into a trivial non-orientable surface-knot by some 1-handle surgeries, [6]. Thus we may assume that F and F' are trivial non-orientable surface-knots with $e(F) = e(F')$. By Whitney's congruence, we have $\chi(F) \equiv \chi(F') \pmod{2}$. Doing 1-handle surgeries if necessary, we may assume that $\chi(F) = \chi(F')$. Then F and F' are ambient isotopic. Thus original 1-component surface-links F and F' are unorientedly bordant. ■

4 Necklaces

We introduce a family of surface-links, called *necklaces*, which is used to prove Theorem 1.2. In the upper 3-space $\mathbf{R}_+^3 = \{(x, y, z) | z \geq 0\}$, we take a 3-ball

$$B^3 = \{(x, y, z) | x^2 + y^2 + (z - 2)^2 \leq 1\}.$$

Let $f = \{f_t\}_{0 \leq t \leq 1}$ and $g = \{g_t\}_{0 \leq t \leq 1}$ be ambient isotopies of B^3 which present a 180° -rotation around the z -axis and a 360° -rotation around the axis $(y = 0, z = 2)$ respectively. We put a Hopf link $k_1 \cup k_2$ in B^3 as in Figure 6 so that $f_1(k_i) = k_i$ and $g_1(k_i) = k_i$ for $i = 1, 2$. By a *motion* of $k_1 \cup k_2$, we mean an ambient isotopy $h = \{h_t\}_{0 \leq t \leq 1}$ of B^3 with $h_1(k_1 \cup k_2) = k_1 \cup k_2$. Two motions h and h' are *equivalent* if there is a 1-parameter family of motions between h and h' .




Figure 6: Spinning the Hopf link in two directions

We consider that $\mathbf{R}^4$ is obtained by spinning $\mathbf{R}_+^3$ around $\partial\mathbf{R}_+^3$ by use of a map $\mu : \mathbf{R}_+^3 \times [0, 1] \rightarrow \mathbf{R}^4$ defined by $((x, y, z) \times \{t\}) \mapsto (x, y, z \cos 2\pi t, z \sin 2\pi t)$. For integers p, q , we construct a 2-component surface-link $S^{p,q} = T_1 \cup T_2$, called a *strand*, as follows:

$$T_i = \mu \left(\bigcup_{t \in [0,1]} (f^p \cdot g^q)_t(k_i) \times \{t\} \right) \subset \mathbf{R}^4 \quad (i = 1, 2).$$

Each T_i ($i = 1, 2$) is homeomorphic to a torus (or a Klein bottle, respectively) if p is even (or odd).

Lemma 4.1 (i) Strands $S^{p,q}$ and $S^{p+2q,0}$ are ambient isotopic.

(ii) If $p \equiv p' \pmod{4}$, then two strands $S^{p,q}$ and $S^{p',q}$ are ambient isotopic.

Proof. By the belt trick (cf. [5]), the motion g is equivalent to f^2 , and f^4 is equivalent to the identity. Hence we have the result. ■

The above lemma implies that ambient isotopy classes of strands are represented by $S^{p,q}$ with $q = 0$ and $p \in \mathbf{Z}_4$. We shall abbreviate $S^{p,0}$ to S^p .

Lemma 4.2 For a strand $S^p = T_1 \cup T_2$, we have

(i) $e(T_1) = e(T_2) = 0 \in \mathbf{Z}$,

(ii) $d(T_1, T_2) = -d(T_2, T_1) = p \in \mathbf{Z}_4$, and

(iii) $t(T_1, T_2, T_1) = t(T_2, T_1, T_2) = p \in \mathbf{Z}_2$.

Proof. (i) For each $i = 1, 2$, we take a 2-disk D_i embedded in B^3 with $\partial D_i = k_i$ and $f_1(D_i) = D_i$. The image $\mu(D_i \times [0, 1])$ is a 3-manifold whose boundary is T_i . Thus $e(T_i) = 0$.

(ii) In Figure 7, we illustrate the motion f of the Hopf link $k_1 \cup k_2$. Since we can obtain a diagram of S^p by taking p copies of the motion and connecting them, we have $d(T_1, T_2) = p$. Similarly, we have $d(T_2, T_1) = -p$.

(iii) The motion in Figure 7 contains two Reidemeister moves of type III. One of them corresponds to a triple point of type (top, middle, bottom) = (T_1, T_2, T_1) and the other corresponds to that of (T_2, T_1, T_2) . Thus we have $t(T_1, T_2, T_1) = t(T_2, T_1, T_2) = p$. ■




Figure 7: The movie of the strand S^1

For m numbers $\{t_i\}_{i=1,\dots,m}$ with $0 < t_1 < \dots < t_m < 1$, we consider a surface-link defined as follows:

$$S^p \cup \mu(\partial B^3 \times \{t_1\}) \cup \dots \cup \mu(\partial B^3 \times \{t_m\}).$$

We call such a surface-link a *necklace* and a spherical component $B_i = \mu(\partial B^3 \times \{t_i\})$ a *bead* of the necklace.

Lemma 4.3 *Let $S^p \cup B_1 \cup \dots \cup B_m$ be a necklace with the strand $S^p = T_1 \cup T_2$. For each $i = 1, \dots, m$, we have*

$$\begin{cases} t(T_2, T_1, B_i) = t(B_i, T_1, T_2) = 1, \\ t(T_1, T_2, B_i) = t(B_i, T_2, T_1) = 1, \\ t(T_1, B_i, T_2) = t(T_2, B_i, T_1) = 0. \end{cases}$$

Proof. In a surface diagram, a bead introduces four triple points $\tau_1, \dots, \tau_4$ as shown in Figure 8. The top, middle and bottom sheets around τ_1 come from T_2 , T_1 and B_i respectively. For τ_2 , they are T_1 , T_2 and B_i . For τ_3 , they are B_i , T_1 and T_2 . For τ_4 , they are B_i , T_2 and T_1 . ■




Figure 8: The local picture of a bead on a strand

We denote by $N^p(i, j; k_1, \dots, k_m)$ an n -component surface-link which is a necklace $S^p \cup B_1 \cup \dots \cup B_m$ with the strand $S^p = T_1 \cup T_2$ and with a label $\alpha : \{T_1, T_2, B_1, \dots, B_m\} \rightarrow \{1, \dots, n\}$ such that $\alpha(T_1) = i$, $\alpha(T_2) = j$, and $\alpha(B_1) = k_1, \dots, \alpha(B_m) = k_m$.

Lemma 4.4 (i) $N^p(i, i; k_1, \dots, k_m)$ is unorientedly null-bordant.

(ii) $N^p(i, j; k_1, \dots, k_m) \cong_A N^{-p}(j, i; k_1, \dots, k_m)$.

(iii) $N^p(i, j; k_1, k_2, k_3, \dots, k_m) \simeq_B N^p(i, j; k_3, \dots, k_m)$ provided $k_1 = k_2$.

(iv) $N^p(i, j; k_1, k_2, \dots, k_m) \simeq_B N^{p+2}(i, j; k_2, \dots, k_m)$ provided $k_1 = i$.

(v) $N^p(i, j; k_1, \dots, k_m) \amalg N^{p'}(i, j; k'_1, \dots, k'_l) \simeq_B N^{p+p'}(i, j; k_1, \dots, k_m, k'_1, \dots, k'_l)$.

(vi) $N^0(i, j; \emptyset)$ is unorientedly null-bordant.

Proof. (i) Consider an annulus A in $\text{int} B^3$ with $\partial A = k_1 \cup k_2$ and $f_1(A) = A$. The image $\mu(A \times [0, 1])$ is a 3-manifold whose boundary is the strand $S^p = T_1 \cup T_2$. Thus, we can remove S^p and then all the beads B_i ($i = 1, \dots, m$) up to unoriented bordism.

(ii) By a 180° -rotation of B^3 around the axis ($y = 0, z = 2$), the components k_1 and k_2 are switched. Then the direction of the rotation f is reversed.

(iii) We can remove the beads $B_i = \mu(\partial B^3 \times \{t_i\})$ ($i = 1, 2$) from F by a 3-manifold $\mu(\partial B^3 \times [t_1, t_2])$.

(iv) We perform a 1-handle surgery between K_1 and B_1 as shown in Figure 9(1) $\rightarrow$ (2). The diagram illustrated in (2) is ambient isotopic to that in (3). This surface is realized by a motion as in (4), which is the 360° -rotation around the axis $(x = 0, z = 2)$. This motion is equivalent to f^2 .

(v) Connect the strands of $N^p(i, j; k_1, \dots, k_m)$ and $N^{p'}(i, j; k'_1, \dots, k'_l)$ up to unoriented bordism to obtain $N^{p+p'}(i, j; k_1, \dots, k_m, k'_1, \dots, k'_l)$ (see Lemma 2.3 of [1]).

(vi) Consider the Hopf link $k_1 \cup k_2$ to use the definition of a strand. Let γ_i ($i = 1, 2$) be disjoint simple arcs in $\mathbf{R}_+^3$ connecting between k_i and $\partial\mathbf{R}_+^3$ in an obvious way. Then the 2-handle surgeries along the 2-disks $\mu(\gamma_1 \times [0, 1])$ and $\mu(\gamma_2 \times [0, 1])$ make S^0 a split union of 2-spheres. ■




Figure 9: The movies of surgeries and isotopies

5 Crossing Changes

In this section, we study a crossing change along a double curve of a surface diagram. The idea is similar to a crossing change for a classical link in $\mathbf{R}^3$. For a classical link, a crossing change can be realized by 1-handle(=band) surgeries as shown in Figure 10(i). Hence, any classical link $k_1 \cup \dots \cup k_n$ is bordant to a split union $k_1 \amalg \dots \amalg k_n \amalg$ (Hopf links).

Lemma 5.1 *Any n -component surface-link $F_1 \cup \dots \cup F_n$ is unorientedly bordant to a split union $F_1 \amalg \dots \amalg F_n \amalg$ (necklaces).*




Figure 10: A movie of a crossing change and the resulting surface

In a surface diagram, we use a symbol E (as in Figure 10(ii)) for a local diagram which is obtained from Figure 10(i) by regarding it a motion picture.

Let c be a double curve of a surface diagram such that it is an immersed loop in $\mathbf{R}^3$. As mentioned in Section 2, a regular neighborhood $N(c)$ of c is regarded as the image of an immersion φ of $B^2 \times [0, 1] / \sim$. See Figure 11. For each c , we choose a pair of diagonal regions Y of $B^2 \setminus X$ and put $R(c) = \varphi(Y \times [0, 1] / \sim)$.




Figure 11: A neighborhood of a double curve

Let A_1 and A_2 be the sheets which intersects along the curve c such that A_1 is higher than A_2 with respect to the projection direction; that is, c is of type (upper, lower) = (A_1, A_2) . To prove Lemma 5.1, we introduce four kinds of local deformations under which unoriented bordism classes are preserved.

(1) Local crossing change. Consider two 1-handle surgeries as shown in Figure 12 in the motion picture method. We call this a *local crossing change* along c . We always assume that the ‘local strand’ $(\text{Hopf link}) \times [0, 1]$ obtained by a local crossing change is in the specified region $R(c)$. Let t be a triple point on c and H a sheet which is transverse to c at t . If H is a top or bottom sheet, then we can perform a local crossing change along the curve c as in Figure 13. For example, if H is a top sheet, then the type of t is changed from (top, middle, bottom) = (H, A_1, A_2) to (H, A_2, A_1) , and the local strand goes under the sheet H .




Figure 12: The local crossing change




Figure 13: A local crossing change near a triple point

(2) End-change. Consider a composite of three ambient isotopies as shown in Figure 14. We call this an *end-change* of a local strand. This deformation moves an end-part of the local strand into the diagonal region of $R(c)$. The new local strand has additional intersections with A_1 and A_2 . In the bottom of Figure 14, the boxed “ f ” means a local diagram corresponding to a 180° -rotation of the Hopf link.




Figure 14: The movies and diagrams determining an end change

(3) Canceling adjacent ends. Assume that two adjacent ends of local strands are in the same region of $R(c)$. Consider a deformation illustrated in Figure 15, which is realized by two 2-handle surgeries on A_1 and A_2 . We call this a *canceling of adjacent ends* of local strands along c .




Figure 15: Canceling adjacent ends for a crossing change

(4) Making a bead. Assume that a strand intersects a sheet H transversely. In case that the strand is under H , we consider a 2-handle surgery on H as shown in Figure 16.

This deformation makes the strand over H with producing a bead.




Figure 16: Making a bead

In a surface diagram of an n -component surface-link $F = F_1 \cup \dots \cup F_n$, a double curve of type (i, j) is *preferred* if $i \leq j$, and a triple point of type (i, j, k) is *preferred* if $i \leq j$ and $j \leq k$.

Proof of Lemma 5.1. Consider a surface diagram of $F = F_1 \cup \dots \cup F_n$. If there is a non-preferred double curve without triple points (that is an embedded loop), we apply a local crossing change along the curve followed by canceling adjacent ends of the local strand. This makes the double curve preferred and yields a strand (necklace without beads), which is separated from F . Thus we may assume that there is at least one triple point on any non-preferred double curve of the surface diagram.

For each non-preferred triple point t of type (i, j, k) , we perform local crossing changes so that t changes into a preferred triple point as follows:

- If $j < i \leq k$, then we perform a local crossing change along the double curve of type (i, j) over the bottom sheet labeled k .
- If $i \leq k < j$, then we perform a local crossing change along the double curve of type (j, k) under the top sheet labeled i .
- If $k < i \leq j$, then we perform a local crossing change along the double curve of type (j, k) and then perform another along the curve of type (i, k) .
- If $j \leq k < i$, then we perform a local crossing change along the double curve of type (i, j) and then perform another along the curve of type (i, k) .
- If $k < j < i$, then we perform a local crossing change along the double curve of type (i, j) , next along the curve of type (i, k) , and then along the curve of type (j, k) .

Figure 17 shows the case of $k < j < i$.

All the local crossing changes that are described above are performed along non-preferred double curves. After applying suitable end-changes, we can cancel all the adjacent ends of local strands along non-preferred double curves. Then we obtain a surface diagram of $(F_1 \amalg \dots \amalg F_n) \cup (\text{strands})$; for any double curves between F_i and F_j are preferred. By making beads if necessary, we can split necklaces from $F_1 \amalg \dots \amalg F_n$. Thus F is unorientably bordant to $F_1 \amalg \dots \amalg F_n \amalg (\text{necklaces})$. ■




Figure 17: A three-fold crossing change at a triple point of type $k < j < i$

Lemma 5.2 $N^0(i, j; k) \simeq_B N^0(k, i; j) \amalg N^0(k, j; i)$.

Proof. Consider a surface diagram of $N^0(i, j; k)$ which contains the local diagram illustrated in Figure 8. In the local diagram, Along the double curves of type (i, k) and (j, k) , we perform (global) crossing changes as in the proof of Lemma 5.1. Then $N^0(i, j; k)$ is unorientably bordant to a split union of $N^0(i, k; j) \cong_A N^0(k, i; j)$ along the curve of type (i, k) , $N^0(j, k; i) \cong_A N^0(k, j; i)$ along the curve of type (j, k) , and a surface-link F obtained by the crossing change. Then F is a split union of $S^0(i, j)$ and a trivial 2-sphere labeled k , which is unorientably null-bordant by Lemma 4.4(vi). ■

6 Proof of Theorem 1.2

Let P_+ and P_- be the standard projective plane in $\mathbf{R}^4$ with $e(P_+) = 2$ and $e(P_-) = -2$ respectively. See Figure 5. We denote by P^m the connected sum of m copies of P_+ if $m > 0$, $-m$ copies of P_- if $m < 0$, and the empty if $m = 0$. Regarding P^m as an n -component surface-link with a label $\alpha(P^m) = i$, we denote it by $P^m(i)$. Regarding a strand $S^p = T_1 \cup T_2$ as an n -component surface-link such that $\alpha(T_1) = i$ and $\alpha(T_2) = j$, we denote it by $S^p(i, j)$. Then $S^p(i, j) = N^p(i, j; \emptyset)$ in the notation used in Lemma 4.4.

Lemma 6.1 Any n -component surface-link F is unorientably bordant to a split union

$$\left(\coprod_{i \in \Gamma_1} P^{m_i}(i) \right) \amalg \left(\coprod_{(i, j) \in \Gamma_2} S^{p_{ij}}(i, j) \right) \amalg \left(\coprod_{(i, j, k) \in \Gamma_3} N^0(i, j; k) \right).$$

Here, Γ_i ($i = 1, 2, 3$) is a subset of the i -fold Cartesian product of $\{1, \dots, n\}$, $m_i \in \mathbf{Z}$ ($i \in \Gamma_1$) and $p_{ij} \in \mathbf{Z}_4$ ($(i, j) \in \Gamma_2$) satisfying the following.

- (i) $m_i \neq 0$ for any $i \in \Gamma_1$.
- (ii) $i < j$ and $p_{ij} \neq 0$ for any $(i, j) \in \Gamma_2$.
- (iii) $i < j < k$ or $i < k < j$ for any $(i, j, k) \in \Gamma_3$.

Proof. By Lemma 5.1, any n -component surface-link $F = F_1 \cup \dots \cup F_n$ is unorientably bordant to $(F_1 \amalg \dots \amalg F_n) \amalg F'$, where F' is a split union of necklaces. Put $\Gamma_1 = \{i \mid e(F_i) \neq 0\}$ and $m_i = e(F_i)/2 \in \mathbf{Z}$ ($i \in \Gamma_1$). By Lemma 3.1, we see that $F_1 \amalg \dots \amalg F_n$ is unorientably bordant to $\coprod_{i \in \Gamma_1} P^{m_i}(i)$ satisfying the condition (i).

Since a necklace $N^p(i, j; k_1, \dots, k_m)$ is unorientedly bordant to $S^p(i, j) \amalg N^0(i, j; k_1) \amalg \dots \amalg N^0(i, j; k_m)$ by Lemma 4.4(v), F' is unorientedly bordant to $F'' \amalg F'''$ such that F'' is a split union of some $S^p(i, j)$'s and F''' is a split union of some $N^0(i, j; k)$'s.

We may assume that $i < j$ for any $S^p(i, j)$ appearing in F'' by Lemma 4.4(i) and (ii). Moreover, by Lemma 4.4(v) and (vi), we see that there exist a subset $\Gamma_2 \subset \{1, \dots, n\}^2$ and $p_{ij} \in \mathbf{Z}_4$ such that F'' is unorientedly bordant to $\coprod_{(i,j) \in \Gamma_2} S^{p_{ij}}(i, j)$ and that the condition (ii) is satisfied.

By Lemma 4.4(i), (ii), (iv) and (vi), we may assume that $i < j$, $i \neq k$ and $j \neq k$ for any $N^0(i, j; k)$ appearing in F''' . Applying Lemma 5.2 for $N^0(i, j; k)$ with $k < i < j$, we may assume that (i, j, k) satisfies the condition (iii) for any $N^0(i, j; k)$ appearing in F''' . By Lemma 4.4(iii), (v) and (vi), we see that there exists a subset $\Gamma_3 \subset \{1, \dots, n\}^3$ such that F''' is unorientedly bordant to $\coprod_{(i,j,k) \in \Gamma_3} N^0(i, j; k)$ and that the condition (iii) is satisfied. ■

For the unoriented bordism group $UL_{4,n}$, we consider three types of homomorphisms e_i ($i = 1, \dots, n$), d_{ij} ($i \neq j$), and t_{ijk} ($i \neq j$ and $j \neq k$) as follows:

$$\begin{cases} e_i : UL_{4,n} \longrightarrow \mathbf{Z} & \text{for } [F] \mapsto e(F_i)/2, \\ d_{ij} : UL_{4,n} \longrightarrow \mathbf{Z}_4 & \text{for } [F] \mapsto d(F_i, F_j), \\ t_{ijk} : UL_{4,n} \longrightarrow \mathbf{Z}_2 & \text{for } [F] \mapsto t(F_i, F_j, F_k), \end{cases}$$

where $F = F_1 \cup \dots \cup F_n$.

Lemma 6.2 *For an n -component surface-link F , let Γ_i ($i = 1, 2, 3$), $m_i \in \mathbf{Z}$ ($i \in \Gamma_1$) and $p_{ij} \in \mathbf{Z}_4$ ($(i, j) \in \Gamma_2$) be as in Lemma 6.1. Then we have the following.*

- (i) $e_i([F]) = m_i$ if $i \in \Gamma_1$ and $e_i([F]) = 0$ if $i \notin \Gamma_1$.
- (ii) For $i < j$, $d_{ij}([F]) = p_{ij}$ if $(i, j) \in \Gamma_2$ and $d_{ij}([F]) = 0$ if $(i, j) \notin \Gamma_2$.
- (iii) For $i < j < k$, $t_{ijk}([F]) = 1$ if $(i, j, k) \in \Gamma_3$ and $t_{ijk}([F]) = 0$ if $(i, j, k) \notin \Gamma_3$.
- (iv) For $i < k < j$, $t_{ijk}([F]) = 1$ if $(i, j, k) \in \Gamma_3$ and $t_{ijk}([F]) = 0$ if $(i, j, k) \notin \Gamma_3$.

Proof. (i) Since $e(P^m) = 2m$, we have $e_i([F]) = m_i$ if $i \in \Gamma_1$ and otherwise $e_i([F]) = 0$.

(ii) This follows from Lemma 4.2(ii).

(iii) and (iv) Note that (j, i, k) , (j, k, i) , (k, i, j) , $(k, j, i) \notin \Gamma_3$. Since $t_{ijk}([N^0(i, j; k)]) = 1$ and $t_{ijk}([N^0(i, k; j)]) = 0$ by Lemma 4.3, $t_{ijk}([F]) = 1$ if and only if $(i, j, k) \in \Gamma_3$. ■

Proof of Theorem 1.2. Consider a homomorphism

$$UH : UL_{4,n} \longrightarrow \underbrace{(\mathbf{Z} \oplus \dots \oplus \mathbf{Z})}_n \oplus \underbrace{(\mathbf{Z}_4 \oplus \dots \oplus \mathbf{Z}_4)}_{\frac{n(n-1)}{2}} \oplus \underbrace{(\mathbf{Z}_2 \oplus \dots \oplus \mathbf{Z}_2)}_{\frac{n(n-1)(n-2)}{3}}$$

defined by $UH = \left(\bigoplus_{i=1}^n e_i \right) \oplus \left(\bigoplus_{i < j} d_{ij} \right) \oplus \left(\bigoplus_{i < j < k \text{ or } i < k < j} t_{ijk} \right)$. This homomorphism is injective by Lemma 6.2. Also, UH is surjective; indeed, $UH([P^1(i)])$ ($i = 1, \dots, n$), $UH([S^1(i, j)])$ ($i < j$), and $UH([N^0(i, j; k)])$ ($i < j < k$ or $i < k < j$) are generators of $\mathbf{Z}$, $\mathbf{Z}_4$, and $\mathbf{Z}_2$ respectively. ■

In the definition of the homomorphism UH , we do not use all double linking numbers and triple linking numbers. The unused ones are determined as follows.

Proposition 6.3 *For distinct i, j, k and an n -component surface-link F , we have*

- (i) $d_{ji}([F]) = -d_{ij}([F])$,
- (ii) $t_{iji}([F]) = t_{jij}([F]) = \lambda(d_{ij}([F]))$, where $\lambda : \mathbf{Z}_4 \rightarrow \mathbf{Z}_2$ is the natural projection,
- (iii) $t_{ijk}([F]) = t_{kji}([F])$, and
- (iv) $t_{jik}([F]) + t_{ijk}([F]) + t_{ikj}([F]) = 0$.

Proof. It is sufficient to prove (i) and (ii) in case that F is as in Lemma 6.1. We use Lemma 4.2. We have $t_{iji}([F]) = p_{ij} \in \mathbf{Z}_2$ if $i < j$, and $t_{iji}([F]) = p_{ji} \in \mathbf{Z}_2$ if $i > j$. On the other hand, we have $d_{ij}([F]) = p_{ij} \in \mathbf{Z}_4$ if $i < j$, and $d_{ij}([F]) = -p_{ji} \in \mathbf{Z}_4$ if $i > j$. Hence, we have $\lambda(d_{ij}([F])) = t_{iji}([F])$. Similarly, since $d_{ij}([F]) = p_{ij} \in \mathbf{Z}_4$ and $d_{ji}([F]) = -p_{ij} \in \mathbf{Z}_4$ for $i < j$, we have $d_{ij}([F]) = -d_{ji}([F])$. (iii) and (iv) are proved in Theorem 3.2 of [11]. ■

We consider the homomorphism $f : L_{4,n} \rightarrow UL_{4,n}$ induced by the map forgetting the orientations of surface-links. For an oriented n -component surface-link F , we can define two kinds of bordism invariants; double linking invariants $D_{ij} : L_{4,n} \rightarrow \mathbf{Z}_2 = \mathbf{Z}/2\mathbf{Z}$ and triple linking invariants $T_{ijk} : L_{4,n} \rightarrow \mathbf{Z}$ (cf. [1]). Then Sanderson's isomorphism

$$H : L_{4,n} \longrightarrow \underbrace{(\mathbf{Z}_2 \oplus \dots \oplus \mathbf{Z}_2)}_{\frac{n(n-1)}{2}} \oplus \underbrace{(\mathbf{Z} \oplus \dots \oplus \mathbf{Z})}_{\frac{n(n-1)(n-2)}{3}}$$

is given by $H = \left(\bigoplus_{i < j} D_{ij} \right) \oplus \left(\bigoplus_{i < j < k \text{ or } i < k < j} T_{ijk} \right)$. From the definitions of these invariants, the forgetful map f is regarded as

$$(\oplus 0) \oplus (\oplus \kappa) \oplus (\oplus \nu) : (\oplus \{0\}) \oplus (\oplus \mathbf{Z}_2) \oplus (\oplus \mathbf{Z}) \rightarrow (\oplus \mathbf{Z}) \oplus (\oplus \mathbf{Z}_4) \oplus (\oplus \mathbf{Z}_2),$$

under the isomorphisms U and UH , where $\kappa : \mathbf{Z}_2 \rightarrow \mathbf{Z}_4$ is the natural inclusion and $\nu : \mathbf{Z} \rightarrow \mathbf{Z}_2$ is the natural projection.

Acknowledgments JSC is being supported by NSF grant DMS-9988107. SK is being supported by Fellowships from the Japan Society for the Promotion of Science. MS is being supported by NSF grant DMS-9988101. The authors would like to thank Uwe Kaiser for providing helpful information.

References

- [1] J.S. Carter, S. Kamada, M. Saito, and S. Satoh, *A theorem of Sanderson on link bordisms in dimension 4*, Algebraic and Geometric Topology, **1** (2001) 299–310.
- [2] J. S. Carter and M. Saito, *Canceling branch points on projections of surfaces in 4-space*, Proc. Amer. Math. Soc. **116** (1992), 229–237.
- [3] ———, *Reidemeister moves for surface isotopies and their interpretations as moves to movies*, J. Knot Theory Ramifications **2** (1993), 251–284.

- [4] ———, *Knotted surfaces and their diagrams*, the American Mathematical Society, Surveys and Monograph Series (Providence 1998).
- [5] L. H. Kauffman, *Knots and physics*, Series on Knots and Everything, 1, World Scientific Publ.
- [6] S. Kamada, *Non-orientable surfaces in 4-space*, Osaka J. Math., **26** (1989), 367–385.
- [7] W. S. Massey, *Proof of a conjecture of Whitney*, Pacific J. Math., **31** (1969), 143–156.
- [8] D. Roseman, *Reidemeister-type moves for surfaces in four dimensional space*, in Banach Center Publications **42** (1998) Knot theory, 347–380.
- [9] B. J. Sanderson, *Bordism of links in codimension 2*, J. London Math. Soc. **35** (1987), 367–376.
- [10] ———, *Triple links in codimension 2*, Colloquia Mathematica Societatis János Bolyai, Topology, Pécs (Hungary), **55** (1989), 457–471.
- [11] S. Satoh, *Triple point invariants of non-orientable surface-links*, Proc. of the first topology meeting of Japan-Mexico (Molelia, July 1999), Topology and its Applications (to appear).
- [12] H. Whitney, *On the topology of differentiable manifolds*, Lectures in topology, Michigan Univ. Press, 1940.